

**DRUG FACTS**

Active ingredient(s)

Purpose

**Use(s)**  
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

**Warnings**

**Do not use**  
if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have**  
liver or kidney disease. Your doctor should determine if you need a different dose.  
an allergic reaction to this product occurs. Seek medical help right away.

**Ask a doctor or pharmacist before use if you are**

**When using this product**  
do not take more than directed. Taking more than directed may cause drowsiness.

**Stop use and ask a doctor if**

**If pregnant or breast-feeding**

**Keep out of reach of children**  
In case of overdose, get medical help or contact a Poison Control Center right away.

<b>Directions</b>	
adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Other information**

- Tamper Evident: do not use if blister units are torn, broken or show any signs of tampering
- store between 20° to 25°C (68° to 77°F)
- protect from excessive moisture

**Inactive ingredients**  
Anhydrous lactose, colloidal silicon dioxide, corn starch, hypromellose, magnesium stearate, microcrystalline cellulose, povidone and sodium lauryl sulfate.